



IONOPTIX MULTICELL: HIGH-THROUGHPUT CALCIUM AND CONTRACTILITY ACQUISITION IN LEFT VENTRICULAR ISOLATED CARDIOMYOCYTES FROM LARGE AND SMALL ANIMALS

Keywords: Calcium, excitation-contraction coupling, isoproterenol

IonOptix System: [MultiCell High Throughput System](#)

Introduction

To date, high-throughput acquisition of isolated cardiomyocyte intracellular calcium ($[Ca^{2+}]_i$) and contractility has been unavailable to investigators seeking to increase throughput. In particular, the acute contractile effects of beta-adrenergic stimulants limit the ability of investigators to generate substantial paired cellular data sets prior to cell death. Beta-adrenergic stimulation accelerates cardiomyocyte excitation-contraction coupling (ECC, for reviews see ¹⁻⁴) kinetics through multiple pathways and can act as a metric of cardiomyocyte functional reserve. Here we show a novel modality for high-throughput, repeated measures acquisition of $[Ca^{2+}]_i$ and contractility prior to and following treatment with the beta adrenergic agonist isoproterenol in enzymatically isolated cardiomyocytes of large and small animals.

Methods

Large Animal Cardiomyocyte Isolation

Cardiomyocytes were isolated from left ventricular midmyocardium of large animal hearts (see ^{5,6} for detailed isolation protocol in porcine model). Following cardiectomy, left ventricular wedge preparations including a portion of the left anterior descending (LAD) coronary artery were excised and cannulated via LAD for coronary perfusion of liberase blendzyme. Ancillary coronary arteries were ligated to improve enzymatic penetration. After sufficient digestion, a portion of the midmyocardium was excised, immersed in bovine serum albumin (BSA) solution to halt enzymatic digestion, and cardiomyocytes were mechanically dissociated from the tissue.

Murine Cardiomyocyte Isolation

Left ventricular cardiomyocytes were isolated from langendorff retrograde perfused hearts of young (3-4 months) mice.⁷⁻¹⁰ Following cardiectomy, hearts were cannulated via the aorta to allow retrograde coronary perfusion of liberase blendzyme. After sufficient digestion, the heart was removed from the cannula and immersed in bovine serum albumin (BSA) solution to halt enzymatic digestion, and cardiomyocytes were mechanically dissociated from the tissue.

Calcium Adaptation and Fura-2/AM Loading

Cardiomyocyte-containing solution was filtered through a 200 μ m filter and slowly adapted to physiologic $[Ca^{2+}]_i$ (2 mM) over 1 hour. During adaptation, cardiomyocytes were loaded with 5 μ M Fura-2/acetoxymethyl ester (Fura-2/AM) and allowed to de-esterify for ratiometric assessment of $[Ca^{2+}]_i$.¹¹

Ratiometric Calcium and Contractility Acquisition

Fura-2 loaded cardiomyocytes were moved to the IonOptix MultiCell stimulation chamber and equilibrated to physiologic temperature (i.e. 35-37° C) for 3 minutes prior to experimentation. Following equilibration, cells were electrically field stimulated at 1.0 Hz, 15 Volts while simultaneously assessing ratiometric $[Ca^{2+}]_i$ (Ex: 340 $[Ca^{2+}$ -bound]/380 $[Ca^{2+}$ -unbound] nm, Em: 510 nm) and contractility (fast fourier transform-derived sarcomere length) at 250 Hz for 10 seconds using the IonWizard software application. Following baseline acquisition under control conditions (CTL), cells were treated with 1 μ M isoproterenol (ISO) to assess cardiomyocyte adrenergic functional kinetic reserve using the MultiCell repeated measures functionality.¹² Increased inotropic effects of isoproterenol resulted in gradual Ca^{2+} overload and spontaneous Ca^{2+} transients, therefore only electrically-stimulated (action-potential induced) transients prior to the advent of spontaneous activity were analyzed.



Results

Multicell Permits Rapid, Paired Measurement of Isolated Cardiomyocyte $[Ca^{2+}]_i$ and Contractility with Isoproterenol Challenge

Using the IonOptix MultiCell High-Throughput repeated measures system, 34 murine (N=1) and 91 large animal (N=2) cardiomyocytes were assessed for systolic and diastolic $[Ca^{2+}]_i$ (**Figure 1 & Figure 3**, respectively) and contractility (**Figure 2 & Figure 4**, respectively) parameters. Data collection from each subject took <1 hour from start to finish and allowed pairwise statistical comparisons using the mark-and-find capability of the MultiCell High Throughput system. Systolic and diastolic parameters and kinetics were augmented in response to isoproterenol perfusion in both species. Unexpectedly, given larger functionally viable cell yields, acquisition times from large animal isolations were significantly shorter than for murine models.

Discussion

The IonOptix Multicell High Throughput System newly empowers investigators to obtain high-throughput calcium and contractility measurements for pairwise comparison in otherwise prohibitive conditions. Given the acute hypercontractility pursuant to beta-adrenergic perfusion, repeated measures assessment of contractility was not always possible, accounting for reduced sample in ISO conditions. However, this did not affect calcium measurements, which were consistent through multiple repeated measures (data not shown). This novel modality allows investigators to expand investigations into cardiomyocyte calcium and contractility beyond the limitations of current modalities.

Bibliography

1. Eisner DA, Caldwell JL, Kistamás K, Trafford AW. Calcium and Excitation-Contraction Coupling in the Heart. *Circ Res* 2017;**121**:181–195.
2. Bers DM. Excitation-Contraction Coupling. 2nd ed. Dordrecht: Kluwer Academic Publishers; 2001.
3. Bers DM. Calcium Cycling and Signaling in Cardiac Myocytes. *Annu Rev Physiol* 2008;**70**:23–49.
4. Bers DM. Cardiac excitation-contraction coupling. *Nature* 2002;**415**:198–205.
5. McDonald KS, Hanft LM, Domeier TL, Emter CA. Length and PKA Dependence of Force Generation and Loaded Shortening in Porcine Cardiac Myocytes. *Biochem Res Int* 2012;**2012**:1–12.
6. Hiemstra JA, Veteto AB, Lambert MD, Olver TD, Ferguson BS, McDonald KS, Emter CA, Domeier TL. Chronic low-intensity exercise attenuates cardiomyocyte contractile dysfunction and impaired adrenergic responsiveness in aortic-banded mini-swine. *J Appl Physiol* 2018;**124**:1034–1044.
7. Liao R, Jain M. Isolation, Culture, and Functional Analysis of Adult Mouse Cardiomyocytes. *Vascular Biology Protocols* 2007. p. 251–262.
8. Louch WE, Sheehan KA, Wolska BM. Methods in cardiomyocyte isolation, culture, and gene transfer. *J Mol Cell Cardiol* 2011;**51**:288–298.
9. Kaestner L, Lipp P, Vecerdeia A, Hammer K, Ruppenthal S, Scholz A. Isolation and Genetic Manipulation of Adult Cardiac Myocytes for Confocal Imaging. *J Vis Exp* 2009;1–5.
10. Veteto AB, Peana D, Lambert MD, McDonald KS, Domeier TL. TRPV4 Contributes to Stretch-Induced Hypercontractility and Time-Dependent Dysfunction in the Aged Heart. *Cardiovasc Res* 2019;1–10.
11. Grynkiewicz G, Poenie M, Tsien RY. A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J Biol Chem* 1985;**260**:3440–3450.
12. Briston SJ, Caldwell JL, Horn MA, Clarke JD, Richards MA, Greensmith DJ, Graham HK, Hall MCS, Eisner DA, Dibb KM, Trafford AW. Impaired β -adrenergic responsiveness accentuates dysfunctional excitation-contraction coupling in an ovine model of tachypacing-induced heart failure. *J Physiol* 2011;**589**:1367–1382.




Figure 1. Isoproterenol Increases $[Ca^{2+}]_i$ and Kinetic Parameters in Murine Cardiomyocytes. (A) Average traces of $[Ca^{2+}]_i$ before (CTL) and after (ISO) 1 μ M isoproterenol perfusion. Isoproterenol increased systolic $[Ca^{2+}]_i$ handling parameters (B-C) and kinetics (D), while augmenting diastolic kinetics (E-G). (N=1, n=34, ****P<0.001 vs. CTL).




Figure 2. Isoproterenol Increases Cardiomyocyte Contractility and Contractile Kinetic Parameters in Murine Cardiomyocytes. (A) Average traces of cardiomyocyte sarcomere length before (CTL) and after (ISO) 1 μ M isoproterenol perfusion. Systolic parameters of contractility showed increased contractility (B-C, F) and kinetic speed (D-E), while relaxation speed was also increased (G). (N=1, n=34, ***P<0.005 vs. CTL, *P<0.05 vs. CTL).




Figure 3. Isoproterenol Increases $[\text{Ca}^{2+}]_i$ and Kinetic Parameters in Large Animal Cardiomyocytes. (A) Average traces of $[\text{Ca}^{2+}]_i$ before (CTL) and after (ISO) $1 \mu\text{M}$ isoproterenol perfusion. Isoproterenol increased systolic $[\text{Ca}^{2+}]_i$ handling parameters (B-C) and kinetics (D), while augmenting diastolic kinetics (E-G). (N=2, n=91, ****P<0.001 vs. CTL).




Figure 4. Isoproterenol Increases Cardiomyocyte Contractility and Contractile Kinetic Parameters in Large Animal Cardiomyocytes. (A) Average traces of cardiomyocyte sarcomere length before (CTL) and after (ISO) 1 μM isoproterenol perfusion. Systolic parameters of contractility showed increased contractility (B-C, F) and kinetic speed (D-E), while relaxation speed was also increased (G). (N=2, n=91, ****P<0.001 vs. CTL, ***P<0.005 vs. CTL).



Adam B. Veteto, Ph.D

Applications Scientist

Post-doctoral Fellow

Office: (617) 696-7335

Mobile: (417) 268-5492

Email: adam@ionoptix.com

Web: www.IonOptix.com